Comparison of the occurrence of tick-borne diseases in ticks collected from vegetation and animals in the same area¹⁾

MONIKA ROCZEŃ-KARCZMARZ, PAULINA DUDKO*, MARTA DEMKOWSKA-KUTRZEPA, MICHAŁ MEISNER**, MARIA STUDZIŃSKA, ANDRZEJ JUNKUSZEW*, ANTONINA Sopińska***, KRZYSZTOF TOMCZUK

Department of Parasitology and Invasive Diseases, Faculty of Veterinary Medicine, University of Life Sciences in Lublin, Akademicka 12, 20-033 Lublin, Poland

*Sub-Department of Small Ruminant Breeding and Professor T. Efner Research Station, Institute of Animal Breeding and Biodiversity Conservation, Faculty of Biology, Animal Science and Bioeconomy, University of Life Sciences in Lublin, Akademicka 13, 20-950 Lublin

**Department of Psychology, Faculty of Social Sciences, The John Paul II Catholic University of Lublin, Al. Racławickie 14, 20-950 Lublin, Poland,

***Department of Fish Diseases and Biology, Faculty of Veterinary Medicine, University of Life Sciences in Lublin, Akademicka 12, 20-033 Lublin, Poland

Received 15.05.2018

Accepted 07.06.2018

Roczeń-Karczmarz M., Dudko P., Demkowska-Kutrzepa M., Meisner M., Studzińska M., Junkuszew A., Sopińska A., Tomczuk K.

Comparison of the occurrence of tick-borne diseases in ticks collected from vegetation and animals in the same area

Summary

The aim of this study was to compare the prevalence of selected pathogens in ticks taken from cats and dogs and from vegetation in urban settlements. A study was conducted to estimate the distribution of *Anaplasma phagocytophilum* and *Borrelia burgdorferi* in adult *Ixodes ricinus* (236), *Dermacentor reticulatus* (237) and *Ixodes hexagonus* (3) ticks collected from animals in veterinary clinics (184) and from vegetation in urban settlements (292). The most numerous ticks collected from animals were *Ixodes ricinus* (73.9%), followed by *Dermacentor reticulatus* (24.5%) and *Ixodes hexagonus* (1.6%). A total of 65.8% of the ticks collected from vegetation were *Dermacentor reticulatus* and 30% were *Ixodes ricinus*. The arthropods removed from the animals were most commonly located around the neck (48.1%) and in the mouth area (17.1%). All ticks were analyzed by molecular techniques. The percentages of ticks positive for *Borrelia burgdorferi* and *Anaplasma phagocytophilum* among those collected from animals differed from the corresponding rates for ticks taken from vegetation in the same area. *Anaplasma phagocytophilum* was more common in ticks collected from vegetation (N = 137, or 47.20%) than in those from animals (N = 12, or 6.6%). *Borrelia burgdorferi*, as well, was more common in ticks collected from the vegetation (N = 96, or 32.9%) than in those from animals (N = 19, or 10.5%). The DNA of *A. phagocytophilum* and *B. burgdorferi* were detected in 30.4% and 22.8% of *D. reticulatus* ticks, respectively, and in 32.6% and 25.4% of *I. ricinus* ticks, respectively. The DNA of *A. phagocytophilum* was also found in one *Ixodes hexagonus* tick. Single infections were noted in 69 *I. ricinus* ticks, 56 *D. reticulatus* ticks and 1 *I. hexagonus* tick. Coinfections of *A. phagocytophilum* with *B. burgdorferi* were detected in 33 (14.0%) *I. ricinus* ticks and in 29 (12.24%) *D. reticulatus* ticks. Infected companion animals can form a reservoir for human tick-transmitted infectious agents. The monitoring of the pathogens transmitted by ticks is an important tool in preventing and combating infections transmitted to humans and animals.

Keywords: *Ixodes ricinus*, *Dermacentor reticulatus*, *Ixodes hexagonus*, *Anaplasma phagocytophilum*, *Borrelia burgdorferi*, pets

¹⁾ The study was conducted under the project „The uses and protection of the genetic resources of farm animals under conditions of sustainable development” co-financed by the National Centre for Research and Development as part of the strategic programme of research and development „Natural environment, agriculture and forestry” – BIOSTRATEG.

Ticks, as hematophages, are a disease transmission vector for both humans and animals. They play an important role in the natural transmission of disease and in transferring pathogens between hosts. The spread of ticks is favoured by a number of ecological, physiological and molecular features. These include high adaptability to changing environmental conditions, climate, the large number of host species, population density in a given area, and the ability to transmit pathogens transovarially (from one generation to the next via the female ovaries) and trans-stadially (from stage to stage: from larvae to nymphs and to adults). The migration of the hosts to geographically distant areas thus aids the spread of ticks and the pathogens transmitted by them (29, 44). The pathogens most often occur in forest ecosystems, in wild animals that constitute a natural source of infection. Ticks are carried away from their natural environment due to the migration of their hosts on which they parasitize, the development of tourism, and frequent trips of the owners of companion animals. In this way, the spread of ticks into new territories contributes to the appearance of new disease entities in areas where they had not previously occurred (42, 36).

In Poland, the most numerous ticks are those from the genus *Ixodes*: *Ixodes ricinus* and *Dermacentor reticulatus*. The former is one of the most important arthropods in the epidemiology of transmission diseases and is considered the most important tick vector in Europe. In Poland, the pathogens most frequently transmitted by *I. ricinus* ticks are *Borrelia burgdorferi* s.l. and *Anaplasma phagocytophilum*, protozoa from the genus *Babesia* and the tick-borne encephalitis virus (KZM) (Flaviviridae) (47, 54, 53). *D. reticulatus* is the second most important vector of many transmissible diseases, and the pathogens most frequently transmitted by these ticks are *Babesia* spp., *Borrelia burgdorferi* s.l., *Anaplasma marginale*, *Rickettsia* spp., and the tick-borne encephalitis virus (KZM) (Flaviviridae) (4, 21, 46). *Ixodes hexagonus* is rarely found in Poland, probably because it is a typically a nest-dwelling species. This tick is involved in the transmission of pathogens that are dangerous to humans and animals, including tick-borne encephalitis, *Borrelia* spp., *Rickettsia* spp., and *Anaplasma phagocytophilum* (23, 29, 40).

Pathogens transmitted by ticks are an important problem in both human and veterinary medicine, with tick-borne diseases being a major threat for humans and animals. The aim of this study was to determine the prevalence of selected pathogens in ticks collected from cats and dogs living in urban habitats and from vegetation in urban settlements.

Material and methods

Study area and tick collection. The study was conducted in south-eastern Poland in the years 2015-2017. The study included 476 ticks (*Ixodes ricinus*, *Dermacentor reticulatus* and *Ixodes hexagonus*), (*Acari*: *Ixodidae*). The ticks were examined in two groups: those collected from the environment and those found on animals brought into the veterinary clinics. Ticks originating from the environment were collected

during the spring and autumn periods of their activity. They were collected by the flagging method from typically urban areas highly frequented by animals and their owners, such as walking routes and city parks. Grass, shrubs and bushes (up to 1.5 m in height) were swept with a white flannel flag (1 × 1 m). The material collected was preserved in plastic sample-tubes. The second group of ticks were collected from cats and dogs (each tick came from a different animal) in several veterinary clinics in Lublin. The parasites were removed from animals with tweezers, described (cat/dog host, age, sex, breed, and location on the host) and placed in an Eppendorf tube. In the laboratory, all ticks were placed in 70% ethyl alcohol. Next, the developmental stage, genus and species of each tick were determined (43) using the Cell light microscope system software from Olympus and then preserved for further molecular studies.

DNA isolation and detection of *Anaplasma phagocytophilum* and *Borrelia burgdorferi*. DNA was isolated from 476 *Ixodes ricinus*, *Dermacentor reticulatus*, and *Ixodes hexagonus* ticks collected from vegetation (292) and from cats and dogs (184). The ticks were removed from ethanol storage and crushed. DNA isolation was performed using a Genomic Mini kit (A&A Biotechnology, Poland), according to the manufacturer's instructions.

Polymerase Chain Reaction (PCR). PCR reactions were carried out to test for the presence of *Borrelia burgdorferi* and *Anaplasma phagocytophilum*. The amplifications were performed using an MJ Research PTC-200 DNA Engine (BioRad, USA). Each PCR reaction was carried out in a 25 µl reaction volume containing 12.5 µl of DreamTaq Green PCR Master Mix (ThermoFisher Scientific, USA), 0.6 µl of 10 µM each of primer (DNA Sequencing and Synthesis Service of the Institute of Biochemistry and Biophysics, Polish Academy of Sciences in Warsaw, Poland.), 3 µl of matrix DNA and 8.3 µl of nuclease-free water supplied for the PCR Master Mix.

Detection of *Borrelia burgdorferi* was carried out using the primers FL6 5'-TTCAGGGTCTCAAGCGTCTTGACT-3' and FL7 5'-GCATTTTCAATTTTAGCAAGTGATG-3', which amplify a product of 276 bp for sequencing the flagellin flaB gene (35). The reactions were performed under the following conditions: initial denaturation at 94°C for 2 minutes, followed by 30 cycles of denaturation at 94°C for 1 minute, annealing at 37°C for 2 minutes, extension at 72°C for 3 minutes, followed by a final extension at 72°C for 3 minutes.

To identify *Anaplasma phagocytophilum*, the following primers were used: EHR 521: 5'-TGTAGGCGGTTC-GTAAGTTAAAG-3' and EHR 747: 5'-GCACTCA-TCGTTTACAGCGT-3', which amplify a product of 274 bp for the 16S rRNA gene (54). The reactions were performed under the following conditions: initial denaturation at 94°C for 5 minutes, followed by 40 cycles with denaturation at 94°C for 45 seconds, annealing at 37°C for 45 seconds, extension at 72°C for 45 seconds, and final extension at 72°C for 5 minutes.

The PCR products were subjected to electrophoresis in 2% agarose gels for the detection of *A. phagocytophilum* and in 1% agarose gels for *B. burgdorferi*; the gels were stained with ethidium bromide and visualized under ultraviolet light.

Statistical analysis. The statistical analysis of parasite occurrence related to the environmental origin involved a chi-square test (including the results of the Z Fisher test with Bonferroni correction for significance level) and two types of nominal correlation depending on the number of categories of the variables (V Cramer and Phi). All analyses were performed using the PS Imago software package (IBM SPSS

Statistics Version 23). The chi-square test and the Z Fisher test were used for precise determination of statistically significant differences between the presence of each parasite associated with an independent variable – the environment. Cramer's phi correlation was used to demonstrate a relationship between the variables and their strength. A p value of ≤ 0.05 was considered significant.

Results and discussion

The detection of pathogens in ticks, as definitive hosts, and in animals, as intermediate hosts, is an important tool in understanding the transmission of pathogens in nature. In this study we analyzed *Ixodes ricinus*, *Dermacentor reticulatus* and *Ixodes hexagonus* ticks for the presence of *Borrelia burgdorferi* and *Anaplasma phagocytophilum*.

Overall, a total of 476 ticks were collected, 184 of which originated from animals (65 from dogs, 119 from cats) and 292 from vegetation. There are statistically significant differences in the numbers of *Dermacentor reticulatus* ($n = 45$) and *Ixodes ricinus* ($n = 136$) collected from the animals ($\chi^2 = 45.75$, $df = 1$, $p < 0.001$). Our studies show that the most common ticks in the animals were *Ixodes ricinus* (73.9%), followed by *Dermacentor reticulatus* (24.5%) and *Ixodes hexagonus* (1.6%). With regard to the ticks collected from vegetation, 65.8% were *Dermacentor reticulatus* and 30% were *Ixodes ricinus*. Most of the ticks removed from animals were located in the neck area (48.1%). Ticks were also removed from the area of the mouth (17.1%), paws (14.9%), abdomen (10.5%) and back (9.4%). The high proportion of ticks found on the head is confirmed by other studies (6).

There are statistically significant differences in the occurrence of *Anaplasma phagocytophilum* ($\chi^2 = 84.99$, $df = 1$, $p < 0.001$) and *Borrelia burgdorferi* ($\chi^2 = 30.41$, $df = 1$, $p < 0.001$) depending on the origin of the ticks. *Anaplasma phagocytophilum* was more common in ticks originating from vegetation ($n = 137$, or 47.20%) than in those from animals ($n = 12$, or 6.6%). *Anaplasma phagocytophilum* was found in one *Ixodes hexagonus* tick, but due to the low numbers of this tick, the results were not taken into account in statistical calculations. *Borrelia burgdorferi* was more common in ticks from the vegetation ($n = 96$, or 32.9%) than in ticks from animals ($n = 19$, or 10.5%).

Single infections were noted in 69 *I. ricinus*, in 56 *D. reticulatus* and in 1 *I. hexagonus* ticks. Coinfections of *Anaplasma phagocytophilum*

with *Borrelia burgdorferi* were detected in 33 (14.0%) *Ixodes ricinus* ticks and in 29 (12.24%) *Dermacentor reticulatus* ticks. The prevalence of *A. phagocytophilum* and *B. burgdorferi* in adult *I. ricinus*, *D. reticulatus* and *I. hexagonus* ticks collected from animals and vegetation is shown in Table 1.

I. ricinus is widespread in Poland, and its numbers are steadily growing (Siuda et al. 1995). The risk of infestation with *D. reticulatus* is also high in the study area, because eastern and central Poland is inhabited by populations of this tick (21, 22). Recreational areas in Poland abound in *I. ricinus* (1, 50, 51, 55), and to a lesser extent in *D. reticulatus* (3, 24, 55). Both *I. ricinus* (10, 12, 20, 49) and *D. reticulatus* ticks (19, 19, 32) can be found in Europe.

In our study, the majority of ticks collected from walking areas in the Lublin districts belonged to the *D. reticulatus* species, while the ticks collected from animals were mostly from the *I. ricinus* species. Very similar results for ticks collected from pets were obtained in other studies in Poland (24) and in Belgium (6). Other authors in Poland (27, 57) found the largest number of *D. reticulatus* ticks on animals, especially dogs. Both species differ in terms of geographic range, seasonal and daily activity peaks and type of habitat (15, 17, 30). Nevertheless, both species occur in most housing estates and walking routes in the Lublin region. Most *I. ricinus* were collected in alleys and housing estates, places where short vegetation passes into dense shrubs

Tab. 1. Prevalence of *Anaplasma phagocytophilum* and *Borrelia burgdorferi* in adult *Ixodes ricinus*, *Dermacentor reticulatus* and *Ixodes hexagonus* ticks collected from animals in urban areas

Location of tick collection	Pathogen	<i>D. reticulatus</i>	<i>I. ricinus</i>	<i>I. hexagonus</i>
Vegetation	<i>B. burgdorferi</i>	51	44	0
	<i>A. phagocytophilum</i>	72	65	0
Cats	<i>B. burgdorferi</i>	2	6	0
	<i>A. phagocytophilum</i>	0	6	0
Dogs	<i>B. burgdorferi</i>	1	10	0
	<i>A. phagocytophilum</i>	0	6	1

Fig. 1. Gel electrophoresis of PCR products in a 2% agarose gel using *Anaplasma phagocytophilum* specific 16S rRNA gene region primers (274 pb PCR product). Lines from 49 to 58 – *D. reticulatus*, and from 59 to 70 – *I. ricinus*

and trees. Also, tall uncropped grass favoured the occurrence of *I. ricinus*. The vicinity of the Bystrzyca River and the Zemborzycki Reservoir favoured the occurrence of *D. reticulatus*. In our study, out of 184 ticks collected from animals, only 3 were *I. hexagonus*. In Europe, ticks of this species are most often found on hedgehogs (8, 11, 34), but they are also found on companion animals (24, 27, 28, 31, 45).

The main vector of *A. phagocytophilum* is the *I. ricinus* tick. The literature mentions few cases of *A. phagocytophilum* transmission by *D. reticulatus* (22, 52, 56). Our study indicates a high risk of *Anaplasma phagocytophilum* and *Borrelia burgdorferi* in the study areas. This study showed high rates of *A. phagocytophilum* infection in *D. reticulatus* and *I. ricinus* ticks. The percentages of *I. ricinus* ticks infected with *A. phagocytophilum* and *B. burgdorferi* were very high: 32.6% and 25.4%, respectively. *Ixodes ricinus* was within the range of infections with *Borrelia* (3.1-40.0%) and *Anaplasma* (1.0-35.0%) in other European countries (5, 6, 10), although in Poland our results indicate the highest level for *Borrelia* (5.6-23.5%) and *Anaplasma* (1.7-19.1%) (9, 13, 24, 39, 48, 51) in urban areas and on walking paths. *Anaplasma phagocytophilum* is found rarely in *D. reticulatus* ticks in Poland and other European countries (1.1-25.36%) (22, 56), whereas *B. burgdorferi* occurs more often and with lower infection rates (0.6-2.7%) (9, 24, 37). In our study, *D. reticulatus* ticks were highly infected with *A. phagocytophilum* (30.4%) and *B. burgdorferi* (22.8%). Our study shows that *A. phagocytophilum* in *D. reticulatus* ticks occurred only in specimens collected from vegetation, and that the rates of infection with both pathogens for *I. ricinus* and *D. reticulatus* ticks were at a similar, high level. This may result from the fact that the primary source of infection occurs in urban settlements, where infected intermediate hosts are likely to occur.

The percentages of ticks collected from animals and positive for *Borrelia burgdorferi* and *Anaplasma phagocytophilum* were 10.5% and 6.6%, respectively. These rates were within the range of infection for *Borrelia* (0.3-22.5%) and *Anaplasma* (1.6-19.5%) found in other European countries (2, 6, 24, 28, 57).

The presence of *B. burgdorferi* and *A. phagocytophilum* in ticks can be determined by the activity of the ticks' hosts inhabiting the biotope, whose occurrence is influenced by a multiple of biotic and abiotic factors (14, 26, 41). The reservoir hosts for *B. burgdorferi* in urban settings are rodents, hedgehogs, foxes, squirrels, hares, birds and deer (14, 16, 25, 38). The typical reservoir hosts for *A. phagocytophilum* are red deer, roe deer, bank voles and wood mice (21, 52). The natural hosts for *I. ricinus*, which are found in forests (Cervidae, Carnivores), most probably also act as reservoirs for *A. phagocytophilum* (51). However, the role of pets should also be taken into account. The development of settlements and their expansion into wasteland, in the vicinity of forests and rivers, increases the access to vectors. Economic and civilizational development often

takes place at the expense of the natural environment. Many species of animals settle and increase in numbers in an uncontrolled manner (7).

There was no significant difference between *I. ricinus* and *D. reticulatus* in the proportion of ticks that contained DNA from both pathogens. All multiple infections were detected in adult *I. ricinus* and *D. reticulatus* ticks, suggesting that they may have fed on the same reservoir hosts. The study of the frequency of coinfection in ticks in a specific area can be used as a tool to help diagnose diseases in humans and animals. In our study, the frequency of coinfection with *A. phagocytophilum* and *B. burgdorferi* was high in both *I. ricinus* and *D. reticulatus*. Other authors have also observed the coincidence of *B. burgdorferi* with *A. phagocytophilum* (33, 54). Such results show that there is a high risk of simultaneous transmission of both pathogens during a single tick bite.

This study shows that there is a high risk of contracting Lyme disease and anaplasmosis in the urban environment, which poses a threat to human and animal populations. It has been shown that, next to *I. ricinus*, *D. reticulatus* is the second most important vector involved in the transfer of *A. phagocytophilum*. Inhabitants and health professionals in cities should be more aware of the risk of Lyme disease and anaplasmosis in green areas or other recreational areas where infected ticks occur.

References

1. Asman M., Solarz K., Cuber P., Gąsior T., Szilman P., Szilman E., Tondaś E., Matzullo A., Kusion N., Florek K.: Detection of protozoans *Babesia microti* and *Toxoplasma gondii* and their co-existence in ticks collected in Tarnogórski district. *Ann. Agric. Environ. Med.* 2015, 22, 50-83, doi: 10.5604/12321966.1141373.
2. Beichel E., Petney T. N., Hassler D., Brückner M., Maiwald M.: Tick infestation patterns and prevalence of *Borrelia burgdorferi* in ticks collected at a veterinary clinic in Germany. *Vet. Parasitol.* 1996, 65, 147-155. 10.1016/0304-4017(96)00943-0.
3. Biernat B., Karbowiak G., Werszko J., Stańczak J.: Prevalence of tick-borne encephalitis virus (TBEV) RNA in *Dermacentor reticulatus* ticks from natural and urban environment, Poland. *Exp. Appl. Acarol.* 2014, 64, 543-551. doi: 10.1007/s10493-014-9836-5.
4. Bonnet S., de la Fuente J., Nicolle P., Liu X., Madani N., Blanchard B., Maingourd G., Alongi A., Torina A., Fernández de Mera I. G., Vicente J., George J. C., Vayssières-Taussat M., Joncœur G.: Prevalence of tick-borne pathogens in adult *Dermacentor* spp. ticks from nine collection sites in France. *Vector-Borne Zoonotic Dis.* 2013, 13, 226-236.
5. Christova I., Van De Pol J., Yazar S.: Identification of *Borrelia burgdorferi* sensu lato, *Anaplasma* and *Ehrlichia* species, and spotted fever group rickettsiae in ticks from Southeastern Europe. *Eur. J. Clin. Microbiol. Infect. Dis.* 2003, 22, 535-542.
6. Claerebout E., Losson B., Cochez C., Casaert S., Dalemans A. C., De Cat A., Madder M., Saegerman C., Heyman P., Lempereur L.: Ticks and associated pathogens collected from dogs and cats in Belgium. *Parasit. Vectors* 2013, 19, 6, 183. doi: 10.1186/1756-3305-6-183.
7. Dudek K., Jerzak L., Tryjanowski P.: Zwierzęta konfliktowe w miastach. Regionalna Dyrekcja Ochrony Środowiska w Gorzowie Wielkopolskim 2016, p. 1-243.
8. Dziemian S., Michalik J., Piłacińska B., Bialik S., Sikora B., Zwolek R.: Infestation of urban populations of the Northern white-breasted hedgehog, *Erinaceus roumanicus*, by *Ixodes* spp. ticks in Poland. *Med. Vet. Entomol.* 2014, 28, 465-469. doi: 10.1111/mve.12065.
9. Dziegiel B., Kubrak T., Adaszek L., Dębiak P., Wylupek D., Bogucka-Kocka A., Lechowski J., Winiarczyk S.: Prevalence of *Babesia canis* *Borrelia burgdorferi* sensu lato and *Anaplasma phagocytophilum* in hard ticks collected from meadows of Lubelskie Voivodship (eastern Poland). *Bull. Vet. Inst. Pulawy* 2014, 58, 29-33.
10. Franke J., Hildebrandt A., Meier F., Straube E., Dorn W.: Prevalence of Lyme disease agents and several emerging pathogens in questing ticks from the German Baltic coast. *J. Med. Entomol.* 2011, 48, 441-444. DOI: 10.1603/ME10182

11. Gern L., Rouvinez E., Toutoungi L. N., Godfroid E.: Transmission cycles of *Borrelia burgdorferi* sensu lato involving *Ixodes ricinus* and/or *I. hexagonus* ticks and the European hedgehog, *Erinaceus europaeus*, in suburban and urban areas in Switzerland. *Folia Parasitol.* 1997, 44, 309-314.
12. Gray J. S., Kirstein F., Robertson J. N., Stein J., Kahl O.: *Borrelia burgdorferi* sensu lato in *Ixodes ricinus* ticks and rodents in a Recreational Park in South-Western Ireland. *Exp. Appl. Acarol.* 1999, 23, 717-729.
13. Grzeszczuk A., Stanczak J., Kubica-Biernat B.: Serological and molecular evidence of human granulocytic ehrlichiosis focus in the Białowieża Primeval Forest, northeastern Poland. *Eur. J. Clin. Microbiol. Infect. Dis.* 2002, 21, 6-11.
14. Grzeszczuk A., Ziarko S., Kovalchuk O., Stańczak J.: Etiology of tick-borne febrile illnesses in adult residents of North-Eastern Poland: report from a prospective clinical study. *Internat. J. Med. Microbiol.* 2006, 296, 242-249.
15. Guglielmone A. A., Robbins R. G., Apanaskevich D. A., Petney T. N., Estrada-Peña A., Horak I.: The hard ticks of the world. Springer, Dordrecht 2014, p. 635.
16. Hamer S. A., Tsao J. I., Walker E. D., Mansfield L. S., Foster E. S., MS, Hickling G. J.: Use of tick surveys and serosurveys to evaluate pet dogs as a sentinel species for emerging Lyme disease. *Am. J. Vet. Res.* 2009, 70, 49-56. doi: 10.2460/ajvr.70.1.49.
17. Hillyard P. D.: Ticks of North-West Europe. The Natural History Museum, London 1996, p. 178.
18. Hornok S., Kartali K., Takács N., Hofmann-Lehmann R.: Uneven seasonal distribution of *Babesia canis* and its two 18S rDNA genotypes in questing Dermacentor reticulatus ticks in urban habitats. *Ticks Tick Borne Dis.* 2016, 7, 694-697. doi: 10.1016/j.ttbdis.2016.02.016.
19. Hornok S., Meli M. L., Gönczi E., Halász E., Takács N., Farkas R., Hofmann-Lehmann R.: Occurrence of ticks and prevalence of *Anaplasma phagocytophilum* and *Borrelia burgdorferi* s.l. in three types of urban biotopes: forests, parks and cemeteries. *Ticks Tick Borne Dis.* 2014, 5, 785-789. doi: 10.1016/j.ttbdis.2014.05.010.
20. Junttila J., Peltomaa M., Soini H., Marjamäki M., Viljanen M. K.: Prevalence of *Borrelia burgdorferi* in *Ixodes ricinus* Ticks in Urban Recreational Areas of Helsinki. *J. Clin. Microbiol.* 1999, 37, 1361-1365.
21. Karbowski G.: Kleszcz łąkowy Dermacentor reticulatus – występowanie, biologia i rola jako wektora chorób odkleszczowych. Rozprawa habilitacyjna. Agencja Reklamowo-Wydawnicza A. Grzegorzczak, Warszawa 2009.
22. Karbowski G., Vichová B., Slivinska K., Werszko J., Didyk J., Pełko B., Stanko M., Akimov I.: The infection of questing Dermacentor reticulatus ticks with *Babesia canis* and *Anaplasma phagocytophilum* in the Chernobyl exclusion zone. *Vet. Parasitol.* 2014, 204, 372-375. doi: 10.1016/j.vetpar.2014.05.030.
23. Krivanec K., Kopecky E., Tomkova E., Grubhoffer L.: Isolation of the TBE virus from the tick *Ixodes hexagonus*, *Folia Parasitol. Praha* 1998, 273-276.
24. Król N., Kiewra D., Szymanowski M., Lonc E.: The role of domestic dogs and cats in the zoonotic cycles of ticks and pathogens. Preliminary studies in the Wrocław Agglomeration (SW Poland). *Vet. Parasitol.* 2015, 30, 214, 208-212. doi: 10.1016/j.vetpar.2015.09.028.
25. Margos G., Vollmer S. A., Cornet M., Garnier M., Fingerle B., Wilske B., Bormane A., Vitorino L., Collares-Periera M., Drancourt M., Kurtenbach K.: MLSA on housekeeping genes defines a new *Borrelia* species. *Appl. Environ. Microbiol.* 2009, 75, 5410-5416.
26. Michalik J., Hofman T., Buczek A., Skoracki M., Sikora B.: *Borrelia burgdorferi* s.l. in *Ixodes ricinus* (Acari: Ixodidae) Ticks Collected from Vegetation and Small Rodents in Recreational Areas of the City of Poznań. *J. Med. Entomol.* 2003, 40, 690-697.
27. Mierzejewska E. J., Welc-Faleciak R., Karbowski G., Kowalec M., Behnke J. M., Bajer A.: Dominance of Dermacentor reticulatus over *Ixodes ricinus* (Ixodidae) on livestock, companion animals and wild ruminants in eastern and central Poland. *Exp. Appl. Acarol.* 2015, 66, 83-101. doi: 10.1007/s10493-015-9889-0.
28. Nijhof A. M., Bodaan C., Postigo M., Nieuwenhuijs H., Opsteegh M., Franssen L., Jebbink F., Jongejan F.: Ticks and associated pathogens collected from domestic animals in the Netherlands. *Vector Borne Zoonotic Dis.* 2007, 7, 1-11. 10.1089/vbz.2007.9999.
29. Nowak-Chmura M.: Fauna kleszczy (Ixodida) Europy Środkowej. Wydawnictwo Naukowe UP, Kraków 2013, s. 212.
30. Nowak-Chmura M., Siuda K.: Ticks of Poland. Review of contemporary issues and latest research. *Ann. Parasitol.* 2012, 58, 125-155.
31. Ogden N. H., Cripps P., Davison C. C., Owen G., Parry J. M., Timms B. J., Forbes A. B.: The ixodid tick species attaching to domestic dogs and cats in Great Britain and Ireland. *Med. Vet. Entomol.* 2000, 14, 332-338. doi: 10.1046/j.1365-2915.2000.00244.x.
32. Olivieri E., Gazzonis A. L., Zanzani S. A., Veronesi F., Manfredi M. T.: Seasonal dynamics of adult Dermacentor reticulatus in a peri-urban park in southern Europe. *Ticks Tick Borne Dis.* 2017, 8, 772-779. doi: 10.1016/j.ttbdis.2017.06.002.
33. Panczuk A., Tokarska-Rodak M., Koziol-Montewka M., Plewik D.: The incidence of *Borrelia burgdorferi*, *Anaplasma phagocytophilum* and *Babesia microti* coinfections among foresters and farmers in eastern Poland. *J. Vector Borne Dis.* 2016, 53, 348-354.
34. Pfäffle M., Petney T., Skuballa J., Taraschewski H.: Comparative population dynamics of a generalist (*Ixodes ricinus*) and specialist tick (*I. hexagonus*) species from European hedgehogs. *Exp. Appl. Acarol.* 2011, 54, 151-164. doi: 10.1007/s10493-011-9432-x.
35. Picken R.: Polymerase chain reaction primers and probes derived from flagellin gene sequences for specific detection of the agents of Lyme disease and North American relapsing fever. *J. Clin. Microbiol.* 1992, 30, 99-114.
36. Ploneczka K., Rypula K., Karczmarczyk R., Szenborn L., Stańczak J.: Badania kleszczy w kierunku zakażenia *Ehrlichia canis* z zastosowaniem reakcji PCR. *Med. Weter.* 2006, 62, 553-556.
37. Reye A. L., Stegny V., Mishaeva N. P., Velhin S., Hübschen J. M., Ignatyev G., Muller C. P.: Prevalence of tick-borne pathogens in *Ixodes ricinus* and Dermacentor reticulatus ticks from different geographical locations in Belarus. *PLOS ONE* 2013, 8, e54476.
38. Rizzoli R., Stevenson J. C., Bauer J. M., van Loon L. J., Walrand S., Kanis J. A., Cooper C., Brandi M. L., Diez-Perez A., Reginster J. Y.: The role of dietary protein and vitamin D in maintaining musculoskeletal health in postmenopausal women: a consensus statement from the European Society for Clinical and Economic Aspects of Osteoporosis and Osteoarthritis (ESCEO). *Maturitas.* 2014, 79, 9, 122-132. doi: 10.1016/j.maturitas.2014.07.005.
39. Roczeń-Karczmarz M.: Zintegrowane działanie pyretroidów na Dermacentor reticulatus (Fabricius, 1794) (Acari: Ixodida: Ixodidae) zebrane we wschodniej Polsce na terenach chronionych o dużym ryzyku zakażenia krętkami *Borrelia burgdorferi* s.l.”. 2009. Doctoral dissertation, Faculty of Medicine with the Dental Department, Medical University of Lublin, Lublin 2009.
40. Schreiber C., Krücken J., Beck S., Maaz D., Pachnicke S., Krieger K., Gross M., Kohn B., von Samson-Himmelstjerna G.: Pathogens in ticks collected from dogs in Berlin/Brandenburg, Germany. *Parasit Vectors.* 2014, 7, doi: 10.1186/s13071-014-0535-1.
41. Siński E., Pawełczyk A., Bajer A., Behnke J.: Abundance of wild rodents, ticks and environmental risk of Lyme borreliosis: a longitudinal study in an area of Mazury Lakes district of Poland. *Ann. Agric. Environ. Med.* 2006, 13, 295-300.
42. Siński E., Welc-Faleciak R.: Risk of infections transmitted by ticks in forest ecosystems of Poland. *Zarządzanie Ochroną Przyrody w Lasach* 2012, tom VI.
43. Siuda K.: Kleszcze Polski (Acari: Ixodidae), cz. II. Systematyka i rozmieszczenie. Monografie Parazytol. nr 12, Wyd. Pol. Tow. Parazyt., Warszawa 1993.
44. Siuda K.: The review of data of the distribution of Ixodida (Acari) in Poland, [in:] Kropczyński D., Boczek J., Tomczyk A. (eds): The Acari, physiological and ecological aspects of Acari-host relationships. Dabor, Warszawa 1995, 273-280.
45. Smith F. D., Ballantyne R., Morgan E. R., Wall R.: Prevalence, distribution and risk associated with tick infestation of dogs in Great Britain. *Med. Vet. Entomol.* 2011, 25, 377-384. doi: 10.1111/j.1365-2915.2011.00954.x.
46. Stańczak J.: Detection of spotted fever group (SFG) rickettsiae in Dermacentor reticulatus (Acari: Ixodidae) in Poland. *Int. J. Med. Microbiol.* 2006, 296, 40, 144-148.
47. Stańczak J., Gabre R. M., Kruminis-Łozowska W., Racewicz M., Kubica B.: *Ixodes ricinus* as a vector of *Borrelia burgdorferi* sensu lato, *Anaplasma phagocytophilum* and *Babesia microti* in urban and suburban forests. *Ann. Agric. Environ. Med.* 2004, 11, 109-114.
48. Stańczak J., Kubica-Biernat B., Racewicz M., Kruminis-Łozowska W., Kur J.: Detection of three genospecies of *Borrelia burgdorferi* sensu lato in *Ixodes ricinus* ticks collected in different regions of Poland. *Int. J. Med. Microbiol.* 2000, 290, 559-566.
49. Svitálková Z., Haruštiaková D., Mahriková L., Berthová L., Slovák M., Kociánová E., Kazimírová M.: *Anaplasma phagocytophilum* prevalence in ticks and rodents in an urban and natural habitat in South-Western Slovakia. *Parasit. Vectors* 2015, 8, 276. doi: 10.1186/s13071-015-0880-8.
50. Wegner Z., Racewicz M., Kubica-Biernat B., Kruminis-Łozowska W., Stańczak J.: Występowanie kleszczy *Ixodes ricinus* (Acari, Ixodidae) na zalesionych obszarach Trójmiasta i ich zakażenia krętkami *Borrelia burgdorferi*. *Przegl. Epidemiol.* 1997, 51, 11-20.
51. Welc-Faleciak R., Kowalec M., Karbowski G., Bajer A., Behnke M. J., Siński E.: Rickettsiaceae and Anaplasmataceae infections in *Ixodes ricinus* ticks from urban and natural forested areas of Poland. *Parasites & Vectors* 2014, 7, 121. DOI: dx.doi.org/10.1186/1756-3305-7-121.
52. Wirtgen M., Nahayo A., Linden A., Garigliany M., Desmecht D.: Detection of *Anaplasma phagocytophilum* in Dermacentor reticulatus ticks. *Vet. Rec.* 2011, 168, 195.
53. Wójcik-Fatla A., Cisak E., Zajac V., Zwoliński J., Dutkiewicz J.: Prevalence of tick-borne encephalitis virus in *Ixodes ricinus* and Dermacentor reticulatus ticks collected from the Lublin region (eastern Poland). *Ticks Tick Borne Dis.* 2011, 2, 16-19. doi: 10.1016/j.ttbdis.2010.10.001.
54. Wójcik-Fatla A., Szmańska J., Wdowiak L., Buczek A., Dutkiewicz J.: Coincidence of three pathogens (*Borrelia burgdorferi* sensu lato, *Anaplasma phagocytophilum* and *Babesia microti*) in *Ixodes ricinus* ticks in the Lublin macroregion. *Ann. Agric. Environ. Med.* 2009, 16, 151-158.
55. Wójcik-Fatla A., Zajac V., Sawczyn A., Cisak E., Dutkiewicz J.: *Babesia* spp. in questing ticks from eastern Poland: prevalence and species diversity. *Parasitol. Res.* 2015, 114, 3111-3116. doi: 10.1007/s00436-015-4529-5.
56. Zajac V., Wójcik-Fatla A., Sawczyn A., Cisak E., Sroka J., Kloc A., Zajac Z., Buczek A., Dutkiewicz J., Bartosik K.: Prevalence of infections and co-infections with 6 pathogens in Dermacentor reticulatus ticks collected in eastern Poland. *Ann. Agric. Environ. Med.* 2017, 24, 26-32. doi: 10.5604/12321966.1233893.
57. Zygmier W., Jaros S., Wędrychowicz H.: Prevalence of *Babesia canis*, *Borrelia afzelii*, and *Anaplasma phagocytophilum* infection in hard ticks removed from dogs in Warsaw (central Poland). *Vet. Parasitol.* 2008, 153, 139-142. doi: 10.1016/j.vetpar.2008.01.036.

Corresponding author: Marta Demkowska-Kutrza, PhD, Akademicka 12, 20-033 Lublin, Poland; e-mail: marta.demkowska@up.lublin.pl